



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

120. Proposed by P. C. CULLEN, Principal of Public Schools, Indianola, Neb.

Draw a circle tangent to a given circle and tangent to a given chord at a given point.

I. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.; G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.; and HENRY HEATON, A. M., Atlantic, Iowa.

Let O be the center of the given circle, and P the given point in the given chord AB .

Through P draw EF perpendicular to chord AB . Draw the diameter COD parallel to EF .

Through C and P draw line CH terminating at H in the circumference of the given circle.

Draw OH intersecting EF at M .

Then will M be the center of a circle tangent to chord AB at P , and tangent to the given circle at H .

PROOF. The center of a circle tangent to AB at P must lie in the perpendicular EF .


The radii OH and MH are drawn to the point of tangency of the two circles. Therefore, the centers O and M and the point of tangency H must lie in the same straight line.

There remains to be proved $MP=MH$.

By construction, MP is parallel to OC , and $OC=OH$. Whence $\triangle COH$ and $\triangle PMH$ are similar.

$\therefore MP=MH$, and M is the center of the required circle, MP and MH being radii thereof.

By a similar construction, we find N the center of a tangent circle on the other side of chord AB , the point of tangency being G .



II. Solution by J. OWEN MAHONEY, B. E., M. Sc., Professor of Mathematics and Science, Cooper Training School, Carthage, Tex.; JOHN J. QUINN, Instructor in Mathematics, Rochester Athenaeum and Mechanics Institute, Rochester, N. Y.; GAYLOR CAMERON, Student Heidelberg University, Tiffin, Ohio; and P. S. BERG, B. Sc., Principal of Schools, Larimore, N. D.

Suppose the problem solved, and let O be the given circle, and E the center of the required circle, and C the given point. One locus of E is the perpendicular to AB at C . From C , on this perpendicular, take $CD=\text{radius of given circle}$; then $EO=ED$. Hence another locus of E is HE , the perpendicular bisector of OE . The intersection of HE and CD determines E .

III. Solution by the PROPOSER.

Let AOB be the given circle, AB the chord, and P the given point, C the center of given circle.

At P erect perpendicular, and with CB as radius construct circle MPN tangent to AB at P . Draw MN intersecting perpendicular at C , which is center of the required circle.

Excellent solutions were received from J. W. YOUNG, J. SCHEFFER, ELMER SCHUYLER, CHAS. C. CROSS, WALTER H. DRANE, ALOIS F. KOVARIK, and P. H. PHILBRICK.